

Exam II

November 5, 2015

Name: _____

This is a 3-hour exam. You may use a calculator and a letter-sized double-sided sheet of notes. No books, cellphones or other electronic devices are allowed. You must show all your work to receive full credit for each problem you solve. The highest score you may receive is 100 points.

1. (10 points) Use a truth table and determine if $(p \wedge q) \rightarrow r$ and $(p \rightarrow r) \wedge (q \rightarrow r)$ are logically equivalent.

2. Let $P(x)$ be the statement " $x^2 > 4x - 2$ " with domain \mathbf{N} .

(a) (5 points) Is $\exists x P(x)$ true or false? Explain.

(b) (5 points) Is $\forall x P(x)$ true or false? Explain.

3. Consider for all real numbers x and y .

(a) (5 points) Is $\forall x \exists y (y = x^2)$ true or false? Explain.

(b) (5 points) Is $\exists y \forall x (y = x^2)$ true or false? Explain.

4. (10 points) Prove directly that *if x and y are rational then $2x + 3y$ is rational.*

5. (10 points) Prove indirectly that *if $3n + 5$ is even then n is odd.*

6. (a) (5 points) Find the first 5 terms for the given sequences.

$$a_0 = a_1 = 1, a_n = a_{n-1} + 2a_{n-2}, n \geq 2.$$

(b) (5 points) Evaluate the summation

$$\sum_{i=5}^8 \left\lfloor \frac{2i + 5}{3} \right\rfloor.$$

7. (10 points) Prove that for all positive integers n ,

$$\sum_{i=0}^n \frac{1}{2^i} = 2 - \frac{1}{2^n}.$$

8. (10 points) Let $P(n)$ be the statement that a gift of n dollars can be formed using just 5-dollar and 6-dollar gift certificates. Prove that $P(n)$ is true for $n \geq 20$.

9. (10 points) Let λ denote the empty binary string. Consider for all possible binary strings made of 0's and 1's. Give a recursive definition for the set S which is all the binary strings that don't have consecutive 0's. For example, $0101101 \in S$ and $110100 \notin S$.
10. (10 points) Give a recursive algorithm to compute the sum of the reciprocals of the first n positive integers. The input to the algorithm is a positive integer n , the output is $\sum_{j=1}^n \frac{1}{j}$.
11. (Bonus Problem.)
- (a) (5 points) Show that $(p \wedge q) \rightarrow r \equiv (p \wedge \neg r) \rightarrow \neg q$.
- (b) (5 points) Use it to prove that *if x is rational and y is irrational then $x + y$ is irrational*.